



## PRINCIPAL COMPONENT ANALYSIS (PCA) OF <sup>1</sup>H NMR AND ANTIFUNGAL ASSAY OF *Hedyosmum brasiliense* MALE AND FEMALE EXTRACTS FROM CERRADO AND ATLANTIC FOREST

**Cynthia Murakami<sup>1</sup>, Jhennifer P. Nastro<sup>2</sup>, Denise M. Selegato<sup>2</sup>, Karina Fraige<sup>2</sup>, Marcos E. L. Lima<sup>1</sup>, Nivaldo Boralle<sup>2</sup>, Vanderlan S. Bolzani<sup>2</sup>, Paulo R. H. Moreno<sup>3</sup>, Maria C. M. Young<sup>4</sup>**

<sup>1</sup>Post Graduation in Plant Biodiversity and Environment, Botanical Institute, São Paulo, Brazil; <sup>2</sup>Nucleus of Bioassays, Biosynthesis and Ecophysiology of Natural Products - NuBBE, - Chemistry Institute, São Paulo State University – UNESP, Araraquara, Brazil; <sup>3</sup>Department of Fundamental Chemistry, Chemistry Institute, University of São Paulo – USP, São Paulo, Brazil; <sup>4</sup>Research Nucleus in Physiology and Biochemistry, Botanical Institute, São Paulo, Brazil; [cynthia.murakami@uol.com.br](mailto:cynthia.murakami@uol.com.br)

**Abstract:** *Hedyosmum brasiliense* is a dioecious shrub from Chloranthaceae, widely distributed in Central, Southeastern and Southern Brazilian regions [1]. Their leaves are popularly used to treat headache, stomach pain, ovarian dysfunction, foot fungi, rheumatism and as diuretic [2]. Since dioecy may influence phytochemical and pharmacological properties of a species [3], so as the environmental conditions where they grow [4], the aim of this work is to investigate the difference among male and female crude extracts of *H. brasiliense* obtained from Cerrado and Atlantic Forest using multivariate analysis and its antifungal activity. Ethanolic extracts were prepared from dried leaves of male and female plants of each biome by cold maceration and 5.0 mg of each extract were lyophilized in triplicate, diluted in 700 µL of CDCl<sub>3</sub>, filtered and transferred to 5 mm glass tubes. A total of 12 samples were analyzed by Hydrogen Nuclear Magnetic Resonance (<sup>1</sup>H NMR, Bruker Ascend 600 MHz). All spectra baselines were adjusted and chemical shifts were referenced by tetramethylsilane (TMS) signals using TopSpin®. Before normalizing the matrix data, regions corresponding to TMS and CDCl<sub>3</sub> of all spectra were suppressed. PCA was executed by NIPALS method using Matlab R2014a Student Version®. For antifungal assay by bioautography, 400 µg of each extract and 5 µg of nystatin (Sigma®) were applied in TLC silica gel 60 F<sup>254</sup> (Merck®), developed with chloroform/acetone (6:1) and registered at (λ) 254 and 366 nm. Suspensions of *Cladosporium cladosporioides* and *C. sphaerospermum* fungi were sprayed separately over the plates, which were incubated for 48 h [5]. Both Atlantic Forest extracts presented strong antifungal activity at Retention Factor (Rf) from 0,53 to 0,67 while those from Cerrado were inactive at the same Rf, suggesting that male and female extracts are chemically similar and the environmental conditions influenced on the biosynthesis of secondary metabolites in this species. However, comparing all extracts by PCA of the <sup>1</sup>H NMR fingerprinting spectrum, it was possible to discriminate Cerrado and Atlantic Forest extracts with 99% of explained variance by PC1xPC2, attributing most of the difference to the signal at δ 1.25 in the NMR spectra. This simple, quick and effective method for crude extract analysis allowed confirming the chemical difference between Cerrado and Atlantic Forest extracts by their antifungal activity and metabolomics.

### References:

- [1] Occhioni, P. 1954. Contribuição ao estudo da família Chloranthaceae com especial referência ao gênero *Hedyosmum* Sw. Tese (Doutorado) - Faculdade Nacional de Farmácia, Rio de Janeiro, 176 p.
- [2] Reitz, R. 1965. Clorantáceas. In: Reitz, R. *et al.* Flora catarinense. Itajaí: Herbário Barbosa Rodrigues, 10 p.
- [3] Simpson, B.S. 2013. Dioecy in plants – Is it an important factor for phytochemists to consider? *Planta Med.* 79: 613-615.
- [4] Karousou, R., Koureas, D.N. and Kokkini, S. 2005. Essential oil composition is related to the natural habitats: *Coridothymus capitatus* and *Satureja thymbra* in NATURA 2000 sites of Crete. *Phytochemistry* 66: 2668-2673.
- [5] Homans, A.L. and Fuchs, A. 1970. Direct bioautography on thin-layer chromatograms as method for detecting fungitoxic substances. *J. Chromatogr.* 51: 327-329.

**Acknowledgements: CNPq and FAPESP.**

---