II CONGRESSO LATINOAMERICANO DE ACAROLOGIA E VI SIMPÓSIO BRASILEIRO DE ACAROLOGIA



29 DE JULHO A O2 DE AGOSTO DE 2018 - PIRENÓPOLIS, GOIÁS, BRASIL ISBN: 978-85-66836-21-9

TAXONOMIC REVISION AND PHYLOGENETIC ANALISYS OF *Tenuipalpus* SENSU STRICTO GROUP (ACARI: TENUIPALPIDAE)

E.B. Castro¹, R. Ochoa², J.J. Beard³, G.R. Bauchan² & R.J.F. Feres¹

¹Depto. Zoologia e Botânica, IBILCE, Universidade Estadual Paulista (UNESP), São José do Rio Preto, SP, Brazil; ²Systematic Entomology Laboratory and Electron and Confocal Microscopy Unit - USDA, ARS, Beltsville, USA; ³Queesland Museum, South Brisbane, Australia.

Tenuipalpus sensu stricto group (Acari: Tenuipalpidae) includes the type species of the genus, T. caudatus (Dugès), and another 39 known species. The definition of this group allowed for a better understanding about of the systematic of the genus, but the relationship among its species is still not well understood. In this study, we provide a taxonomic revision of the Tenuipalpus sensu stricto group based mainly on study of type specimens deposited in the NMNH Mite Collection and in the MCZ Mite Collection. The taxonomic revision allowed the construction of a character matrix in order to propose a phylogenetic analysis for this group of mites. The ingroup consisted of 27 species of *Tenuipalpus* sensu stricto. The outgroup was composed by two species of the genus Ultratenuipalpus. The dataset consisted of 60 morphological characters. These characters were based mainly from gnathosoma, dorsal, ventral and leg chaetotaxy. The analysis identified two equally parsimonious trees of 218 steps (CI: 32, RI: 54). In the strict consensus tree, a branch including the ingroup was supported by eight synapomorphies, including the presence of lateral body projections associated with setae c3. The first major branch was supported by two synapomorphies, related to presence of ridges from prodorsal setae scl to the sejugal furrow. The second major branch was supported by two synapomorphies, including the presence of ventral setae 3a2. Our preliminary results show the presence of two major groups within Tenuipalpus sensu stricto. The inclusion of more species with new characters will collaborate a better understanding about the relationship among the species of this group of mites.

Keywords: flat mites, false spider mites, taxonomy, systematic, body projections, *Ultratenuipalpus*.

Financial support: FAPESP, CNPq.